



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A NEW SUBSPECIES OF HUDSONIAN CHICKADEE
FROM THE LABRADOR PENINSULA.

BY CHARLES W. TOWNSEND, M.D.

***Penthestes hudsonicus nigricans*, subsp. nov.**

Subsp. Char.—Midway in size between *Penthestes hudsonicus hudsonicus* and *P. hudsonicus littoralis*, bill as short as that of *littoralis* and as thick as *hudsonicus*. Darker than either on back and sides and much less brown.

Type.—No. 1420, coll. C. W. T. ♂ ad. Shekatika, head of inlet Saguenay County, Quebec, Canadian Labrador, July 23, 1915; collected by C. W. Townsend.¹

Range.—So far as known, forested region of Labrador Peninsula.

Description of adult. Sexes alike. Crown, nape and back mouse-gray slightly tinged with hair brown; wings and tail dull slate-color; throat black, ear coverts, sides of neck, breast and belly white; sides brownish drab.

Measurements ♂ (type) wing 66, tail 64, tarsus 15, bill, culmen, 8.5, depth at base 4.5.

♀ wing 62, tail 62, tarsus 14, bill, culmen, 8, depth at base 5.

Remarks. The short, stout bill and dark back almost devoid of brown tint as well as the absence of a strong brown tint on the sides make this a well marked subspecies. In its dark sooty tendency it resembles other birds of the forested regions of the Labrador Peninsula such as the Labrador Horned Owl and the Labrador Jay. This region abounds in water in the bogs, lakes and rivers and its climate in summer is humid. A number of other birds of this region appear to have a tendency to darkness in plumage, *e. g.* Red-tailed Hawk, Flicker, Night Hawk and Water-Thrush.

It is probable that the Hudsonian Chickadees of the treeless or scrub Arctic and Subarctic area in Ungava Labrador and Newfoundland Labrador are *P. hudsonicus hudsonicus*. Specimens from Lance au Loup taken in May belong to this latter form. The Ungava form described by Rhoads in 1893 (Auk, X, 1893, p. 328) has the brown coloration of *hudsonicus*. I am greatly indebted to Mr. Outram Bangs for his assistance in preparing this note.

¹ I have since given the type specimen to the Museum of Comparative Zoölogy at Cambridge, Mass., and it now bears the number 69431.